

M.K. HOME TUITION

Mathematics Revision Guides

Level: AS / A Level

AQA : C4

Edexcel: C4

OCR: C4

OCR MEI: C4

THE BINOMIAL SERIES FOR RATIONAL POWERS

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 + \dots + x^n$$

$$\frac{1}{1+x} = (1+x)^{-1} = 1 + (-1)x + \frac{(-1)(-2)}{2!} x^2 + \frac{(-1)(-2)(-3)}{3!} x^3 + \frac{(-1)(-2)(-3)(-4)}{4!} x^4 + \dots$$

$$= 1 - x + x^2 - x^3 + x^4 \dots \quad |x| < 1$$

$$x = 0.05 \quad \frac{1}{1.05} = 1 - 0.05 + 0.0025 - 0.000125 + 0.00000625 = 0.952381$$

$$\frac{1}{\sqrt{4+x}} = (4+x)^{-\frac{1}{2}} = 4^{-\frac{1}{2}} \left[\left(1 + \frac{x}{4} \right)^{-\frac{1}{2}} \right] = \frac{1}{2} \left[\left(1 + \frac{x}{4} \right)^{-\frac{1}{2}} \right]$$

$$= \frac{1}{2} \left[1 + \left(-\frac{1}{2}\right) \left(\frac{x}{4}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!} \left(\frac{x}{4}\right)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!} \left(\frac{x}{4}\right)^3 + \dots \right]$$

$$= \frac{1}{2} - \frac{1}{16} x + \frac{3}{256} x^2 - \frac{5}{2048} x^3 + \dots \quad \left| \frac{x}{4} \right| < 1, \text{ or } |x| < 4.$$

$$\frac{1}{(1-4x)^2} = (1-4x)^{-2}$$

$$= 1 + (-2)(-4x) + \frac{(-2)(-3)}{2!} (-4x)^2 + \frac{(-2)(-3)(-4)}{3!} (-4x)^3 + \dots$$

$$= 1 + 8x + 48x^2 + 256x^3 + \dots$$

$$(1-x)^n = 1 - nx + \frac{n(n-1)}{2!} x^2 - \frac{n(n-1)(n-2)}{3!} x^3 + \dots + x^n$$

THE BINOMIAL SERIES FOR RATIONAL n .

To recap, the general binomial expansion for $(a + b)^n$, where n is a positive integer, is

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1}b + \binom{n}{2} a^{n-2}b^2 + \binom{n}{3} a^{n-3}b^3 + \dots + b^n$$

or

$$(a + b)^n = a^n + na^{n-1}b + \frac{n(n-1)}{2!} a^{n-2}b^2 + \frac{n(n-1)(n-2)}{3!} a^{n-3}b^3 + \dots + b^n.$$

Notice the pattern here: powers of a decrease from n to zero, powers of b increase from zero to n , and in each term there is a binomial coefficient multiplier with a factorial denominator and a product of a series of numbers in the numerator.

Recall factorials: $1! = 1$; $2! = 2 \times 1 = 2$; $3! = 3 \times 2 \times 1 = 6$; $4! = 4 \times 3 \times 2 \times 1 = 24$ and so forth.

Substituting $(1 + x)^n$ for $(a + b)^n$ we have

$$(1 + x)^n = 1 + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots + x^n$$

or

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \frac{n(n-1)(n-2)}{3!} x^3 + \dots + x^n$$

This expression is simpler than the previous one, because all powers of 1 are equal to 1 itself.

Finally, substituting $(1 - x)^n$ for $(1 + x)^n$ in the last expression we have

$$(1 - x)^n = 1 - \binom{n}{1}x + \binom{n}{2}x^2 - \binom{n}{3}x^3 + \dots + (-x)^n$$

or

$$(1 - x)^n = 1 - nx + \frac{n(n-1)}{2!} x^2 - \frac{n(n-1)(n-2)}{3!} x^3 + \dots + (-x)^n$$

The terms in odd powers of x have their sign reversed – think of $(1 - x)$ as $(1 + (-x))$.

If n is a positive integer, the series will terminate at the x^n term, since the numerator of the fractional representation of the binomial coefficient will have a zero term in it, and as multiplying by 0 gives 0, this term and all subsequent ones will vanish.

However, the formulae for $(1 + x)^n$ and $(1 - x)^n$ can also be used for all other rational n , giving an infinite series. Provided x lies within certain limits, the series will converge, in other words, the terms will become smaller as we move from left to right.

A series of the form $(1 + x)^n$ converges, i.e. the expansion is valid, when $|x| < 1$.

If the second term of the binomial is kx where k is a non-zero constant, the limits of convergence are $|kx| < 1$, or $|x| < \frac{1}{k}$.

Example(1): Expand $\frac{1}{1+x}$ up to the term in x^4 and state the values for which the expansion is valid.

Use the result to find the value of $\frac{1}{1.05}$ to 6 decimal places.

The expression can be written as $(1+x)^{-1}$ and therefore its binomial expansion is

$$(1+x)^{-1} = 1 + (-1)x + \frac{(-1)(-2)}{2!} x^2 + \frac{(-1)(-2)(-3)}{3!} x^3 + \frac{(-1)(-2)(-3)(-4)}{4!} x^4 + \dots$$
$$= 1 - x + x^2 - x^3 + x^4.$$

This expansion is valid and convergent for $|x| < 1$.

When $x = 0.05$, the series sums to $1 - 0.05 + 0.0025 - 0.000125 + 0.00000625 = 0.952381$ to 6 d.p.

Note that for examination purposes, questions will usually be restricted only to the term in x^2 or x^3 .

Example(2): Expand $\frac{1}{(1+2x)^2}$ up to the term in x^3 and state the values for which the expansion is valid.

Use the result to estimate $\frac{1}{(1.06)^2}$ to 4 decimal places.

The expression can be written as $(1+2x)^{-2}$ and therefore its binomial expansion is

$$1 + (-2)(2x) + \frac{(-2)(-3)}{2!} (2x)^2 + \frac{(-2)(-3)(-4)}{3!} (2x)^3 + \dots$$

$$\text{or } 1 - 4x + 12x^2 - 32x^3 \dots$$

This expansion is valid and convergent for $|2x| < 1$, or $|x| < \frac{1}{2}$.

When $1 + 2x = 1.06$, $2x = 0.06$, and therefore $x = 0.03$.

Substituting $x = 0.03$ in the expansion, we have $1 - 4(0.03) + 12(0.03)^2 - 32(0.03)^3 \dots$
or $1 - 0.12 + 0.0108 - 0.000864 = 0.8899$ to 4 d.p.

Sometimes, we may need algebraic manipulation to put the expression 'into shape', i.e as some multiple of $(1+x)^n$ or $(1-x)^n$.

Example(3): Expand $\sqrt{1-x}$ up to the term in x^3 , stating the values for which the expansion is valid.

Use the result to find the value of $\sqrt{24}$ to 5 decimal places. (Hint : $24 = 25 \times 0.96$)

The expression can be written as $(1-x)^{\frac{1}{2}}$ and therefore its binomial expansion is

$$(1-x)^{\frac{1}{2}} = 1 - \frac{1}{2}x + \frac{(\frac{1}{2})(-\frac{1}{2})}{2!}x^2 - \frac{(\frac{1}{2})(-\frac{1}{2})(-\frac{3}{2})}{3!}x^3 + \dots \quad \text{or treat as}$$

$$(1-x)^{\frac{1}{2}} = 1 + \frac{1}{2}(-x) + \frac{(\frac{1}{2})(-\frac{1}{2})}{2!}(-x)^2 + \frac{(\frac{1}{2})(-\frac{1}{2})(-\frac{3}{2})}{3!}(-x)^3 + \dots$$

$$= 1 - \frac{1}{2}x - \frac{1}{8}x^2 - \frac{1}{16}x^3 \dots$$

(The binomial coefficients are more awkward here because of the fractions)

This expansion is valid and convergent for $|x| < 1$.

To find $\sqrt{24}$, we cannot substitute $x = -23$ and say $\sqrt{1-(-23)}$, as the series is only valid for $|x| < 1$.

We can, though, manipulate surds to give $\sqrt{24} = \sqrt{25 \times 0.96} = \sqrt{25}\sqrt{0.96} = 5\sqrt{0.96}$.

Now we can calculate $\sqrt{0.96}$ by substituting $x = 0.04$ (to give $1-x = 0.96$) into the binomial series, since x now is within the valid range for expansion.

$$\sqrt{0.96} \approx 1 - \frac{1}{2}(0.04) - \frac{1}{8}(0.0016) - \frac{1}{16}(0.000064)$$

$$\approx 1 - 0.02 - 0.0002 - 0.000004 \approx 0.979796.$$

Multiplying the result by 5 gives $\sqrt{24} = 4.89898$ to 5 decimal places.

Example(4): Expand $\frac{1}{\sqrt{4+x}}$ up to the term in x^3 , stating the values for which the expansion is valid.

Firstly, $\frac{1}{\sqrt{4+x}}$ is the same as $(4+x)^{-\frac{1}{2}}$

The expression inside the square root sign is not of the correct format $(1+x)$ for substituting into the binomial expansion, so we have to take out a factor of 4 as follows :

$$\begin{aligned} \text{Because } (4+x) &= 4\left(1+\frac{x}{4}\right), \text{ we can say } (4+x)^{-\frac{1}{2}} = \left(4\left(1+\frac{x}{4}\right)\right)^{-\frac{1}{2}} \\ &= 4^{-\frac{1}{2}} \left[\left(1+\frac{x}{4}\right)^{-\frac{1}{2}}\right] \text{ (recall laws of indices !)} = \frac{1}{2} \left[\left(1+\frac{x}{4}\right)^{-\frac{1}{2}}\right]. \end{aligned}$$

We now have the all-important 1 as the first term, so we can carry out the expansion :

$$\begin{aligned} \frac{1}{2} \left[\left(1+\frac{x}{4}\right)^{-\frac{1}{2}}\right] &= \frac{1}{2} \left[1 + \left(-\frac{1}{2}\right)\left(\frac{x}{4}\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}\left(\frac{x}{4}\right)^2 + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{3!}\left(\frac{x}{4}\right)^3 \dots\dots\right] \\ &= \frac{1}{2} - \frac{1}{16}x + \frac{3}{256}x^2 - \frac{5}{2048}x^3 \dots\dots \text{ (Do not forget the external factor of } \frac{1}{2} \text{).} \end{aligned}$$

This expansion is valid and convergent for $|\frac{x}{4}| < 1$, or $|x| < 4$.

Example (5):

i) Expand $\frac{1}{(1-4x)^2}$ in ascending powers of x , up to and including the term in x^3 .

ii) Find the coefficient of x^2 in the expansion of $\frac{(1+3x)^2}{(1-4x)^2}$.

$$\text{i) } \frac{1}{(1-4x)^2} = (1-4x)^{-2}$$

$$= 1 + (-2)(-4x) + \frac{(-2)(-3)}{2!}(-4x)^2 + \frac{(-2)(-3)(-4)}{3!}(-4x)^3 + \dots$$

$$= 1 + 8x + 48x^2 + 256x^3 \dots$$

ii) Multiplying the expansion in i) by $(1+3x)^2$ gives the result

$$(1 + 8x + 48x^2 + 256x^3 \dots)(1 + 6x + 9x^2).$$

The combinations of terms contributing to the quadratic term in the product are $(1 \times 9x^2)$, $(8x \times 6x)$ and $(48x^2 \times 1)$. Their sum is $(9 + 48 + 48)x^2$ or $105x^2$.

\therefore the coefficient of x^2 in the expansion of $\frac{(1+3x)^2}{(1-4x)^2}$ is 105.

Example (6): In the section “Partial Fractions”, we resolved the expression

$$\frac{4x+5}{(x-1)(x+2)^2} \text{ into partial fractions as } \frac{1}{x-1} - \frac{1}{x+2} + \frac{1}{(x+2)^2}.$$

An examination question could continue as follows:

i) Use the formula for the sum to infinity of a geometric series to show that the binomial expansion of $\frac{1}{x-1}$ forms the series $-1 - x - x^2 - x^3 \dots$

ii) Show that the expression $\frac{1}{x+2}$ is equivalent to $\frac{1}{2}\left(1 + \frac{x}{2}\right)^{-1}$ and hence find the binomial expansion of $\frac{1}{x+2}$ up to and including the term in x^3 .

iii) Using similar reasoning to part ii), find the binomial expansion of $\frac{1}{(x+2)^2}$ up to and including the term in x^3 .

iv) Hence show that the binomial expansion (to the term in x^3) of $\frac{4x+5}{(x-1)(x+2)^2}$ can be expressed as $-\frac{1}{16}(20 + 16x + 15x^2 + 17x^3)$

v) Hence use the series from iv) to find the value of $\frac{4x+5}{(x-1)(x+2)^2}$ when $x = 0.01$.

vi) State the range of values of x for which the expansion is valid.

i) The series $-1 - x - x^2 - x^3 \dots$ (**Series A**) is recognisable as a G.P. whose first term a is -1 and whose common ratio r is x . Its sum to infinity is therefore $\frac{a}{1-r}$ or in this case $\frac{-1}{1-x}$ or $\frac{1}{x-1}$.

This expansion is valid for $|x| < 1$.

ii) The expression $\frac{1}{x+2}$ cannot be expanded in the format $(2+x)^{-1}$, so we must rewrite it as

$$(2^{-1})\left(1 + \frac{x}{2}\right)^{-1} \text{ and hence as } \frac{1}{2}\left(1 + \frac{x}{2}\right)^{-1}.$$

This expansion is valid for $|\frac{x}{2}| < 1$, or $|x| < 2$.

Expanding up to the term in x^3 we have:

$$\begin{aligned} \frac{1}{2}\left(1 + \frac{x}{2}\right)^{-1} &= \frac{1}{2}\left[1 + (-1)\left(\frac{x}{2}\right) + \frac{(-1)(-2)}{2!}\left(\frac{x}{2}\right)^2 + \frac{(-1)(-2)(-3)}{3!}\left(\frac{x}{2}\right)^3 + \dots\right] \\ &= \frac{1}{2}\left[1 - \frac{1}{2}x + \frac{1}{4}x^2 - \frac{1}{8}x^3 + \dots\right] = \frac{1}{2} - \frac{1}{4}x + \frac{1}{8}x^2 - \frac{1}{16}x^3 \dots \text{ (Series B)} \end{aligned}$$

iii) The binomial expansion of $\frac{1}{(x+2)^2}$ must similarly be adjusted as follows:

$$\begin{aligned} \frac{1}{(x+2)^2} &= (2^{-2})\left(1 + \frac{x}{2}\right)^{-2} = \frac{1}{4}\left(1 + \frac{x}{2}\right)^{-2} \\ &= \frac{1}{4}\left[1 + (-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \frac{(-2)(-3)(-4)}{3!}\left(\frac{x}{2}\right)^3 + \dots\right] \\ &= \frac{1}{4}\left[1 - x + \frac{3}{4}x^2 - \frac{1}{2}x^3 + \dots\right] = \frac{1}{4} - \frac{1}{4}x + \frac{3}{16}x^2 - \frac{1}{8}x^3 \dots \text{ (Series C)} \end{aligned}$$

This expansion is valid for $|\frac{x}{2}| < 1$, or $|x| < 2$.

iv) Combining the expressions for Series **A**, **B** and **C** gives the following results:
 (Remember that series **B** must be subtracted, not added !)

Series A	-1	-x	-x ²	-x ³
Series B (SUBTRACT !)	$\frac{1}{2}$	$-\frac{1}{4}x$	$\frac{1}{8}x^2$	$-\frac{1}{16}x^3$
Series C	$\frac{1}{4}$	$-\frac{1}{4}x$	$\frac{3}{16}x^2$	$-\frac{1}{8}x^3$
Total (A - B + C)	$-\frac{5}{4}$	-x	$-\frac{15}{16}x^2$	$-\frac{17}{16}x^3$

v) The binomial expansion (to the term in x^3) of $\frac{4x+5}{(x-1)(x+2)^2}$ is therefore

$$-\left(\frac{5}{4}\right) - x - \left(\frac{15}{16}x^2\right) - \left(\frac{17}{16}x^3\right) \dots$$

By multiplying everything out by -16 and putting $-\frac{1}{16}$ outside the brackets, the expansion can also be written as $-\frac{1}{16}(20 + 16x + 15x^2 + 17x^3)$.

When $x = 0.01$, $\frac{4x+5}{(x-1)(x+2)^2} \approx -\frac{1}{16}(20 + 0.16 + 0.0015 + 0.000017)$ or 1.260095.

Note: The actual value is $\frac{5.04}{(-0.99)(2.01)^2} = -1.2600948$.

vi) Two of the terms in the combined expression are valid for $|\frac{x}{2}| < 1$, or $|x| < 2$. However, the expression for $\frac{1}{x-1}$ is valid only for $|x| < 1$.

The series *as a whole* is therefore valid only for the range of its 'strictest' component, i.e. $|x| < 1$.