

## M.K. HOME TUITION

Mathematics Revision Guides  
Level: AS / A Level

AQA : C4

Edexcel: C4

OCR: C4

OCR MEI: C4

## PARAMETRIC DIFFERENTIATION

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx} \Rightarrow \frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$
$$x = 8t, y = 2t^3$$
$$\frac{dx}{dt} = 8 \quad \frac{dy}{dt} = 6t^2 \Rightarrow \frac{dy}{dx} = \frac{6t^2}{8} \Rightarrow \frac{dy}{dx} = \frac{3t^2}{4}$$
  
$$x = 3t^2, y = 2t^3 + 6t \Rightarrow \frac{dx}{dt} = 6t, \frac{dy}{dt} = 6t^2 + 6$$
$$\Rightarrow \frac{dy}{dx} = \frac{dy/dt}{dx/dt}, \frac{dy}{dx} = \frac{6t^2 + 6}{6t} \Rightarrow \frac{dy}{dx} = t + \frac{1}{t} \quad \therefore \text{When } t = 5, \frac{dy}{dx} = 5\frac{1}{5}$$
  
$$x = 2 \sin \theta, y = \cos \theta \Rightarrow \frac{dx}{d\theta} = 2 \cos \theta, \frac{dy}{d\theta} = -\sin \theta$$
$$\Rightarrow \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta}, \frac{dy}{dx} = \frac{-\sin \theta}{2 \cos \theta} \Rightarrow \frac{dy}{dx} = -\frac{1}{2} \tan \theta \quad \therefore \text{When } \theta = \pi/4, \frac{dy}{dx} = -\frac{1}{2}$$
  
$$x = t^3, y = 5t^2 - 10$$
$$\Rightarrow \frac{dx}{dt} = 3t^2, \frac{dy}{dt} = 10t \Rightarrow \frac{dy}{dx} = \frac{10t}{3t^2} \Rightarrow \frac{dy}{dx} = \frac{10}{3t}$$

When  $x = 8, t = 2, y = 10, \frac{dy}{dx} = \frac{5}{3} \Rightarrow$  gradient of normal to  $(8, 10) = -\frac{3}{5}$   
equation of normal to  $(8, 10)$  is  $y - 10 = -\frac{3}{5}(x - 8)$   
 $\Rightarrow 5y - 50 = -3x + 24 \Rightarrow 3x + 5y - 74 = 0$

## Parametric Differentiation.

If  $x$  and  $y$  are expressed in terms of another variable  $t$  (a **parameter**), then by the chain rule,

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx} \text{ which can be rearranged as } \frac{dy}{dx} = \frac{dy/dt}{dx/dt}, \text{ since } \frac{dx}{dt} = \frac{1}{dt/dx}.$$

**Example (1):** A curve is defined parametrically by  $x = 3t^2$ ,  $y = 2t^3 + 6t$ .

Find  $\frac{dy}{dx}$  in terms of  $t$ , and give its value when  $t = 5$ .

Differentiating with respect to  $t$  gives  $\frac{dx}{dt} = 6t$  and  $\frac{dy}{dt} = 6t^2 + 6$ .

$$\text{Using } \frac{dy}{dx} = \frac{dy/dt}{dx/dt}, \frac{dy}{dx} = \frac{6t^2 + 6}{6t} \Rightarrow \frac{dy}{dx} = t + \frac{1}{t}.$$

$$\therefore \text{When } t = 5, \frac{dy}{dx} = 5\frac{1}{5}.$$

**Example (2):** A curve is defined parametrically by  $x = 2 \sin \theta$ ,  $y = \cos \theta$ .

(Trigonometric functions use  $\theta$  as a parameter rather than  $t$ ).

Find  $\frac{dy}{dx}$  in terms of  $\theta$ , and give its value when  $\theta = \pi/4$ .

Differentiating with respect to  $\theta$  gives  $\frac{dx}{d\theta} = 2 \cos \theta$  and  $\frac{dy}{d\theta} = -\sin \theta$ .

$$\text{Using } \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta}, \frac{dy}{dx} = \frac{-\sin \theta}{2 \cos \theta} \Rightarrow \frac{dy}{dx} = -\frac{1}{2} \tan \theta.$$

$$\therefore \text{When } \theta = \pi/4, \frac{dy}{dx} = -\frac{1}{2}.$$

**Example (3):** The parametric equation of a curve is  $x = \cos \theta$ ,  $y = 3 \cos 2\theta$ ,  $0 < \theta < \pi$ .

Find  $\frac{dy}{dx}$  in terms of  $\theta$ , and hence find the maximum value that the gradient can take at any point on the curve.

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} \Rightarrow \frac{dy}{dx} = \frac{-6 \sin 2\theta}{-\sin \theta} \Rightarrow \frac{dy}{dx} = \frac{12 \sin \theta \cos \theta}{\sin \theta} \Rightarrow \frac{dy}{dx} = 12 \cos \theta$$

(We can cancel  $\sin \theta$  out as a factor in the last step, since it is never zero when  $0 < \theta < \pi$ ).

Since  $\cos \theta$  takes a maximum value of 1, the gradient must take a maximum value of 12.

Examination questions usually include revision of AS-level techniques such as maxima, minima, tangents and normals.

**Example (4):** A curve is defined parametrically by  $x = 8t$ ,  $y = 2t^3$ .

Find  $\frac{dy}{dx}$  in terms of  $t$  and therefore find the equation of the tangent at the point (16, 16).

Differentiating with respect to  $t$  gives  $\frac{dx}{dt} = 8$  and  $\frac{dy}{dt} = 6t^2$ .

$$\Rightarrow \frac{dy}{dx} = \frac{6t^2}{8} \Rightarrow \frac{dy}{dx} = \frac{3t^2}{4} \text{ by the chain rule.}$$

When  $y = 16$ ,  $t = 2$ , therefore the gradient of the tangent to the point (16, 16) is 3 and its equation is  $y - 16 = 3(x - 16)$ ,  $\Rightarrow y = 3x - 32$  or  $3x - 48 - y + 16 = 0 \Rightarrow 3x - y - 32 = 0$ .

**Example (5):** A curve is defined parametrically by  $x = t^3$ ,  $y = 5t^2 - 10$ .

Find  $\frac{dy}{dx}$  in terms of  $t$  and therefore find the equation of the normal at the point (8, 10).

Differentiating with respect to  $t$  gives  $\frac{dx}{dt} = 3t^2$  and  $\frac{dy}{dt} = 10t$ .

$$\Rightarrow \frac{dy}{dx} = \frac{10t}{3t^2} \Rightarrow \frac{dy}{dx} = \frac{10}{3t}$$

When  $x = 8$ ,  $t = 2$ , and so the gradient of the tangent is  $\frac{5}{3}$ .

The gradient of the normal to the point (8, 10) is thus  $-\frac{3}{5}$ .

Its equation is therefore  $y - 10 = -\frac{3}{5}(x - 8)$

$$\Rightarrow 5y - 50 = -3x + 24 \Rightarrow 3x + 5y - 74 = 0,$$

$$\text{or } y = \frac{74}{5} - \frac{3}{5}x.$$

### Parametric Second Derivatives (Not all syllabuses)

**Example (6):** Extension to Example (4)

A curve is defined parametrically by  $x = 8t$ ,  $y = 2t^3$ .

(We need the following results from Example (3):  $\frac{dy}{dx} = \frac{3t^2}{4}$ ,  $\frac{dx}{dt} = 6t^2$ ).

Find the value of  $\frac{d^2y}{dx^2}$  when  $t = 4$ .

To find the second derivative **with respect to  $x$** , we must use the chain rule again.

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d}{dx} \left( \frac{3t^2}{4} \right) \\ &= \frac{d}{dt} \left( \frac{3t^2}{4} \right) \times \frac{dt}{dx}\end{aligned}$$

**This is important** – we cannot just differentiate with respect to  $t$  and leave it as  $\frac{3t}{2}$ .

Given  $\frac{dx}{dt} = 6t^2$  from Example (5), it follows that  $\frac{dt}{dx} = \frac{1}{6t^2}$ .

Therefore  $\frac{d^2y}{dx^2} = \frac{3t}{2} \times \frac{1}{6t^2} = \frac{3t}{12t^2} = \frac{1}{4t}$ .

When  $t = 4$ ,  $\frac{d^2y}{dx^2} = \frac{1}{16}$ .