

## M.K. HOME TUITION

Mathematics Revision Guides  
Level: GCSE Higher Tier

# SEQUENCES

The diagram illustrates three sequences:

- Square Numbers:** Shown as a grid of squares. For  $n=1$ , there is 1 square. For  $n=2$ , there is a 2x2 grid (4 squares). For  $n=3$ , there is a 3x3 grid (9 squares). For  $n=4$ , there is a 4x4 grid (16 squares). The sequence of terms is  $1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$ .
- Triangular Numbers:** Shown as a triangle of squares. For  $n=1$ , there is 1 square. For  $n=2$ , there are 3 squares. For  $n=3$ , there are 6 squares. For  $n=4$ , there are 10 squares. The sequence of terms is  $1, 5, 9, 13, 17, 21, \dots$ .
- Powers of 2:** Shown as a triangle of triangles. For  $n=1$ , there is 1 triangle. For  $n=2$ , there are 3 triangles. For  $n=3$ , there are 6 triangles. For  $n=4$ , there are 10 triangles. The sequence of terms is  $1, 2, 6, 24, 120, 720, 5040, \dots$ .

Additional sequences shown are  $2, 4, 8, 16, 32, 64, \dots$  and  $1, 2, 6, 24, 120, 720, 5040, \dots$ .


## SEQUENCES

A **sequence** is a list of numbers following some rule for finding succeeding values.

Each number in a sequence is called a **term**.

A sequence can be defined by a formula for the  $n^{\text{th}}$  term, where  $n$  is the position of the term in the sequence.

**Example (1):** One example of a sequence is that of the square numbers, illustrated below.




The first term of the sequence is 1, the second 4, the third 9 and the fourth, 16.

The number of squares in each of the diagrams is the same as the square of its position in the sequence.

The 5<sup>th</sup> term in the sequence will therefore be  $5^2$  or 25, and we can also generalise by saying that the  $n^{\text{th}}$  term is equal to  $n^2$ .

**Example (2):** Investigate the sequence of rectangles below to find a general formula for the width, the height, and thus the area, in unit squares, of the  $n^{\text{th}}$  rectangle.



The first rectangle is 2 squares wide  $\times$  1 square high.

The second rectangle is 3 squares wide  $\times$  2 squares high.

The third rectangle is 4 squares wide  $\times$  3 squares high.

The fourth rectangle is 5 squares wide  $\times$  4 squares high.


We can see an obvious pattern here.

The height of each rectangle equals the position in the sequence – it can be represented as  $n$ .


The width of each rectangle exceeds the position in the sequence by 1 – it can be represented as  $n + 1$ .

Since the area of a rectangle is given by height  $\times$  width, the area of the  $n^{\text{th}}$  one is  $n(n + 1)$ .


**Example (3):** Investigate the sequence of T-shapes below to draw the next T-shape in the sequence, and also to find a general formula for the total number of unit squares in the  $n^{\text{th}}$  T-shape.



$n = 1$



$n = 2$




$n = 3$

Spotting the pattern here is not quite as easy as in the last example, but we can see that:

An extra unit square is added to the vertical bar of the T as  $n$  goes up by 1.

Two extra unit squares (one on each side) are added to the horizontal bar of the T as  $n$  goes up by 1.

The fourth T-shape in the sequence therefore looks like the diagram on the right.



The first T-shape is made up of 5 unit squares.

The second one is made up of  $5 + (3 \times 1)$  or 8 unit squares.

The third one is made up of  $5 + (3 \times 2)$  or 11 unit squares.

The fourth one is made up of  $5 + (3 \times 3)$  or 14 unit squares.

We can spot the pattern here: the  $n^{\text{th}}$  T-shape is made up of  $5 + 3(n-1)$  unit squares.

(An equally valid expression is  $2 + 3n$ ).

The first three examples were shape-based and intuitive, but it is also possible to determine the general term of a sequence by listing enough of its elements to spot a pattern.

**Sequences with a common difference.**

These sequences are generated by **adding a constant value** to each term as we go along.

We can see that the next two terms in the sequence 7, 14, 21, 28 .... are 35 and 42, the terms being the first multiples of 7. We can also see that the  $n^{\text{th}}$  term is  $7n$ .

Since each term can be obtained from the previous one by adding 7, the common difference is 7.

The common difference can be found by taking a term after the first, and subtracting the one before it.

The general formula for the  $n^{\text{th}}$  term is  $u_n = an + b$  where  $a$  is the common difference and  $b$  is a constant to be added. This constant is a hypothetical “0<sup>th</sup> term” derived by subtracting the common difference from the first term.

Note the use of the subscripted variable  $u_n$  to denote the  $n^{\text{th}}$  term of a sequence. By this same notation, the first three terms of a sequence are  $u_1, u_2$  and  $u_3$ .

(This type of sequence is also called an **arithmetic progression**.)

**Examples (4a):** Find the next two terms of the following sequences, together with a general formula for the  $n^{\text{th}}$  term:

- i) 1, 5, 9, 13, 17, .....
- ii) 44, 39, 34, 29, 24, .....

In i) we see that the first term is equal to 1 and the common difference is 4.

The next two terms are 21 and 25.

The formula for the general term is therefore  $u_n = 4n + b$ .

The first term is 1, so the “0<sup>th</sup> term”,  $u_0$ , is  $1 - 4$  or  $-3$ , giving a general formula of  $u_n = 4n - 3$ .

The 100<sup>th</sup> term,  $u_{100}$ , is therefore  $(4 \times 100) - 3$  or 397.

In ii) we can see that successive terms decrease in value, so the common difference will be negative.

The first term  $u_1 = 44$  and the common difference is  $-5$ .

The next two terms are 19 and 14.

This time the expression for the general term is  $u_n = -5n + b$ .

The first term is 44, so  $u_0 = 44 - (-5)$  or 49, giving a general formula of  $u_n = 49 - 5n$ .

Hence  $u_{100} = 49 - (5 \times 100)$  or  $-451$ .

**Example (4b):** For the sequence 1, 5, 9, 13, 17, .....

- i) Find the position of the number 61.
- ii) Show that the number 75 is not a term of the sequence.

i) Because the formula  $u_n = 4n - 3$  can be used to find the  $n^{\text{th}}$  term in the sequence, we solve the equation  $4n - 3 = 61$ , and the solution is  $n = 16$ .

The number 61 is therefore the 16<sup>th</sup> term in the sequence.

ii) When we solve  $4n - 3 = 75$  as in i), we have the result  $n = 19.5$ , which is not a positive whole number, so 75 is not a member of the sequence.

**Sequences with a common ratio.**

These sequences are generated by **multiplying** each term by a **constant value** as we go along.

Take the sequence 3, 9, 27, 81 .... as an example.

The common ratio can be found by taking a term after the first, and dividing it by the one before it.

The general expression for the  $n^{\text{th}}$  term is  $b(a^n)$  where  $a$  is the common ratio and  $b$  is a constant multiplier. This constant is a hypothetical “0<sup>th</sup> term” obtained by dividing the first term by the common ratio.

Since each term can be obtained from the previous one by multiplying by 3, the common ratio is 3.

We can see that the next two terms in the sequence 3, 9, 27, 81 .... are 243 and 729, the terms being the first powers of 3. We can also see that the  $n^{\text{th}}$  term,  $u_n$ , is  $3^n$ .

(This type of sequence is also called a **geometric progression**.)

**Examples (5):** Find the next two terms of the following sequences, together with a general expression for the  $n^{\text{th}}$  term:

- i) 3, 6, 12, 24, 48, .....
- ii) 800, 80, 8, 0.8, 0.08,.....

In i) we see that the first term,  $u_1$ , is equal to 3 and the common ratio  $r$  (e.g.  $6 \div 3$ ) is 2.

The next two terms are 96 and 192.

The first term,  $u_1$  is 3, so  $u_0 = 3 \div 2$  or 1.5, with the resulting general formula  $u_n = 1.5(2^n)$ .

In ii) the successive terms decrease in value, and so the common ratio will be less than 1.

The first term is 800 and the common ratio is 0.1.


The next two terms are 0.008 and 0.0008.

Since the first term,  $u_1$ , is 800, the “0<sup>th</sup> term”,  $u_0$  is  $800 \div 0.1$  or 8000, with the resulting formula of  $u_n = 8000(0.1^n)$  for the general  $n^{\text{th}}$  term.

### Quadratic Sequences (For 2017)

These are trickier than the last two, because we have two sets of differences to consider.

**Example (6):** Take the sequence of triangular numbers.



The triangular numbers are represented here by the shaded triangles.

The first triangular number is 1.

The second triangular number is  $1 + 2$  or 3.

The third triangular number is  $1 + 2 + 3$  or 6.

The fourth triangular number is  $1 + 2 + 3 + 4$  or 10.

The sequence of triangular numbers therefore goes 1, 3, 6, 10, .....

The difference between the first term and the second is 1.

The difference between the second term and the third is 2.

The difference between the third term and the fourth is 3.

The first differences form an arithmetic sequence, but the second differences are constant at 1.

All quadratic sequences have a constant second difference.

We can set up a list of differences as follows :

Sequence terms	1	3	6	10
1 <sup>st</sup> differences		2	3	4
2 <sup>nd</sup> differences		1	1	

To find the next two terms of the series, we put two more instances of 1 in the “2<sup>nd</sup> difference” row.

Then we can put  $4 + 1 = 5$  and  $5 + 1 = 6$  into the “1<sup>st</sup> differences” row above.

Finally, we can put  $10 + 5 = 15$  and  $15 + 6 = 21$  into the top row to get the next triangular numbers.

Sequence terms	1	3	6	10	15	21
1 <sup>st</sup> differences		2	3	4	5	6
2 <sup>nd</sup> differences		1	1	1	1	1

Sequence terms	1	3	6	10	15
1 <sup>st</sup> differences		2	3	4	5
2 <sup>nd</sup> differences		1	1	1	

This method is not suitable if we wanted to find, say, the 60<sup>th</sup> triangular number, since filling in such a long list would be tedious. We need to find the expression for a general term for the sequence,  $u_n$ .

A general term of this sequence is  $u_n = an^2 + bn + c$  where  $a$ ,  $b$  and  $c$  are constants.

We can find  $a$  by halving the second difference.

In the case of the triangular numbers, the second differences are all equal to 1, so  $a = \frac{1}{2}$ .

Hence  $u_n = \frac{1}{2}n^2 + bn + c$ .

Position $n$	1	2	3	4	5
Sequence term $u_n$	1	3	6	10	15
Value of $\frac{1}{2}n^2$	$\frac{1}{2}$	2	$4\frac{1}{2}$	8	$12\frac{1}{2}$
$bn + c$ (after subtracting)	$\frac{1}{2}$	<b>1</b>	<b><math>1\frac{1}{2}</math></b>	<b>2</b>	<b><math>2\frac{1}{2}</math></b>

After subtracting  $\frac{1}{2}n^2$  from each term, we are left with a linear sequence  $u_n = bn + c$  with a common difference of  $\frac{1}{2}$  and a first term of  $\frac{1}{2}$ .

The general term of this linear sequence is  $u_n = \frac{1}{2}n$ , and so the general term of the original is

$$u_n = \frac{1}{2}n^2 + \frac{1}{2}n \text{ or } \frac{1}{2}n(n + 1).$$

Hence the 60<sup>th</sup> triangular number is  $\frac{1}{2} \times 60 \times 61 = 1830$ .

The result resembles that of the rectangle example (2).

Each rectangle can be split into two triangles as shown to the right.


The illustrated example with the triangular numbers could have been solved by other intuitive methods, but sometimes the above method is the only option.




**n = 1**



**n = 2**



**n = 3**



**n = 4**

**Example (7):**

- i) Find the next two terms of the sequence 4, 7, 16, 31, 52.....
- ii) Find the general term for the  $n$ th term  $u_n$ .
- iii) Hence find the 10<sup>th</sup> term.

i) Calculate the differences first :

Sequence terms	4	7	16	31	52	<b>79</b>	<b>112</b>
1 <sup>st</sup> differences	3	9	15	21	27	<b>33</b>	
2 <sup>nd</sup> differences		6	6	6	6	6	

The first differences for the sequence 3, 9, 15, 21 .... and the second differences are constant at 6.

We can therefore put two more instances of 6 in the “2<sup>nd</sup> difference” row.  
 Then we can put  $21 + 6 = 27$  and  $27 + 6 = 33$  into the “1<sup>st</sup> differences” row above.  
 Finally, we put  $52 + 27 = 79$  and  $79 + 33 = 112$  into the uppermost row.

The next two terms of the sequence are therefore 79 and 112.

ii) The general term is  $u_n = an^2 + bn + c$ .

The second differences are all equal to 6, and half of that is 3, so  $a = 3$  and  $u_n = 3n^2 + bn + c$ .

Position $n$	1	2	3	4	5
Sequence term $u_n$	4	7	16	31	52
Value of $3n^2$	3	12	27	48	75
$bn + c$ (after subtracting)	<b>1</b>	<b>-5</b>	<b>-11</b>	<b>-17</b>	<b>-23</b>

After subtracting  $3n^2$  from each term, we are left with a linear sequence  $u_n = bn + c$  with a common difference of -6 and a first term of 1.

The general formula for the  $n$ <sup>th</sup> term of this linear sequence is  $u_n = -6n + 7$ , hence the general  $n$ <sup>th</sup> term of the original sequence is  $u_n = 3n^2 - 6n + 7$ .

iii) We find the 10<sup>th</sup> term of the sequence by substituting  $n = 10$  into the formula  $u_n = 3n^2 - 6n + 7$ .

Thus  $u_{10} = 300 - 60 + 7 = \mathbf{247}$ .



### Generating sequences by rules.

Another way of generating a sequence is to follow some rule relating the terms.  
Thus the sequence 1, 5, 9, 13, 17... can be generated by the rules:

- Start with 1.
- Generate the next term by adding 4 to the term before it.

Another way of expressing these rules is to use an **inductive definition** using subscripted variables :

$$u_1 = 1 ; \quad u_{k+1} = u_k + 4$$

Thus  $u_1$  is the first term of the sequence ( here 1),  $u_2$  the second, and  $u_k$  the general  $k$ th term.

The statement  $u_{k+1} = u_k + 4$  means “To find the next term, take the current one and add 4.”

Similarly the sequence 3, 6, 12, 24, 48, .... can be generated by the rules:

- Start with 3.
- Generate the next term by doubling the term before it

In inductive definition form:

$$u_1 = 3 ; \quad u_{k+1} = 2u_k$$

Thus  $u_1$  , the first term of the sequence, is 3.

Also, the statement  $u_{k+1} = 2u_k$  means “To find the next term, double the current one.”

One disadvantage of this definition is that it is not always easy to deduce a formula for the  $n^{\text{th}}$  term.

### Example (8):

Generate the first ten terms of the sequence generated by the following rule:

- Set the first two terms to 1 and 1.
- Generate each further term by adding together the two previous ones.

Also, write out an inductive definition for the sequence.

(Do not attempt to find a general formula for the  $n^{\text{th}}$  term !)

The third term is 1 + 1 or 2, the fourth one is 1 + 2 or 3 and the fifth one is 2 + 3 or 5.

The sequence thus goes 1, 1, 2, 3, 5, 8, 13, 21, 34, 55 .....

The inductive definition is  $u_1 = 1 ; \quad u_2 = 1 ; \quad u_{k+1} = u_{k-1} + u_k$

Aside: This is the ‘Fibonacci’ sequence, named after a 13<sup>th</sup> – Century mathematician, Leonardo of Pisa, ‘Fibonacci – son of Bonaccio’ who first investigated it.