


M.K. HOME TUITION

Mathematics Revision Guides
Level: GCSE Higher Tier

RECOGNISING GRAPHS OF FUNCTIONS





RECOGNISING GRAPHS



Graphs of functions fall into various categories, but most of those you'll come across will be of a few basic types.

Straight-line graphs (recalled).







Constant graphs:




The main diagonals:



Other straight line graphs passing through the origin:



General straight line graphs, not passing through the origin:



The point where a linear graph cuts the y -axis is also known as the y -intercept, or simply the intercept.

The point where the graph cuts the x -axis is sometimes called the x -intercept, but is more often called the root (as in the solution of an equation).

The more common form of equation of a straight-line graph encountered at GCSE is the form

$y = mx + c$, known as the **gradient-intercept** form.

The graph of $y = 2x - 3$ has a gradient of 2 and a y -intercept at $(0, -3)$.

Similarly the graph of $y = 1 - x$ had a gradient of -1 and its y -intercept was the point $(0, 1)$.

(Note that $y = 1 - x$ is the same as $y = -x + 1$).





The gradient of a graph whose equation is given in gradient-intercept form is evidently the multiple of x , and that the graph crosses the y -axis where y takes the value of the constant.

- Any graph of the form $y = mx + c$ has a gradient of m and a y -intercept at $(0, c)$.

(The gradient-intercept form cannot be applied to constant graphs of the form $x = c$, since they are vertical lines and attempting to find a gradient would mean dividing by zero, which is undefined.)

Quadratic graphs.

These graphs are of functions of the form $y = ax^2 + bx + c$ where a , b and c are constants, and a is not zero. The highest power of x is 2 (the square of x). The basic graph of $y = x^2$ is shown upper left.



These graphs are parabolic or 'bucket-shaped'.

When the x^2 term is positive, the graphs point downwards at a trough and the function takes a minimum value. The expansion of $y = (x - 2)(x - 8)$ is $y = x^2 - 10x + 16$.

On the other hand, they point upwards at a crest and have a maximum value when the x^2 term is negative. The expansion of $y = (1 - x)(2 + x)$ is $y = 2 - x - x^2$.





The 'depth' of a parabolic graph can vary, but this is as dependent on the scaling of the graph axes as on the actual function.

Cubic graphs.

These are a little more complicated than quadratic graphs.

Their functions are of the form $y = ax^3 + bx^2 + cx + d$ where a, b, c and d are constants, and a is not zero. The highest power of x is 3 (the cube of x).

The basic graph of $y = x^3$ is shown upper left.





These graphs are characterised by a 'double bend'.

If the term in x^3 is positive, the general slope is upward from lower left, but if the x^3 term is negative, the general slope is downward from upper left.

The 'bend' can also vary in severity - the graph on lower left has sharper 'bends' than the one on lower right.

Reciprocal graphs.



These graphs are shared by functions of the form $y = \frac{k}{x}$, where k is a non-zero constant. They differ from previous examples in that they seem to be in two unconnected parts: if k is positive, the two sections are in the upper right and lower left, but if k is negative, the sections are in the upper left and lower right.

Note the following features of the standard graph $y = \frac{1}{x}$.

As x becomes large and positive, y stays positive but approaches zero.

As x becomes large and negative, y stays negative but approaches zero.

As positive x approaches zero, y becomes increasingly large and positive.

In other words, y **tends to infinity** (∞).


As negative x approaches zero, y becomes increasingly large and negative,

i.e. y tends to minus infinity ($-\infty$).

When $x = 0$, y is undefined - you cannot divide by zero, so it is absurd to say that $\frac{1}{0} = \infty$.

Exponential Graphs. (A+)

These are graphs of $y = a^x$ where a is any *positive* number.



Graphs of this form have the following features in common:

Regardless of the value of a , $y = 1$ when $x = 0$.

For $x > 1$:

As x increases, y also increases for all values of x .

As x become large and negative, y tends to zero but never gets there.

For $0 < x < 1$:

y would still be 1 when $x = 0$.



As x decreases, y also increases for all values of x .

As x becomes large and positive, y tends to zero but never gets there.

(The graph of $y = a^x$ when $a = 1$ is merely the straight line $y = 1$.)


Trigonometric Graphs. (A+)

The three main trigonometric functions have the following graphs:



The graphs of $\sin x^\circ$ and $\cos x^\circ$ are similar to each other; both functions can only take values in the range -1 to $+1$, and both repeat themselves every 360° . Indeed, the graph of $\cos x^\circ$ is the same as that of $\sin x^\circ$ translated 90° to the left.


The graph of $\tan x^\circ$ is quite different. It repeats every 180° , and moreover the function is undefined for certain values of x , such as 90° , 270° , and all angles consisting of an odd number of right angles. When x approaches 90° from below, $\tan x^\circ$ becomes very large and positive; when x approaches 90° from above, $\tan x^\circ$ becomes very large and negative.



Circular Graphs. (From 2017)


A circle passing through the origin has an equation of $x^2 + y^2 = r^2$, where r is the radius of the circle.

In the example shown on the right, the circle has a radius of 5 units. Remember that the number on the right-hand side of the equation is the *square* of the radius, and not the radius itself.



Equation of the tangent to a circle.

For any point (a,b) on a circle centred on the origin, and with a radius r , the equation of the tangent to the circle at that point is $ax + by = r^2$.



In the above example, the equation of the circle is $x^2 + y^2 = 25$, and the selected point on its circumference is (4, 3).

The equation of the tangent is thus $4x + 3y = 25$.